

Discrete Toda Field Equations.

R. S. WARD

*Isaac Newton Institute for Mathematical Sciences,
Cambridge CB3 0EH, UK*

and

*Dept of Mathematical Sciences, University of Durham,
Durham DH1 3LE, UK.**

Abstract. There are two-dimensional Toda field equations corresponding to each (finite or affine) Lie algebra. The question addressed in this note is whether there exist integrable discrete versions of these. It is shown that for certain algebras (such as A_n , $A_n^{(1)}$ and B_n) there do, but some of these systems are defined on the half-plane rather than the full two-dimensional lattice.

To appear in Physics Letters A.

* Permanent address.

The original Toda chain

$$\ddot{\varphi}_j = \exp - \sum_k C_{jk} \varphi_k, \quad (1)$$

where the C_{jk} are certain constants, is an integrable system in which $\varphi_j(t)$ depends on a discrete variable j and on a continuous variable t [1]. This can be generalized in two ways, while maintaining integrability. First, the number of continuous variables can be increased from one to two (replace t by u, v), with the second derivative being replaced by the two-dimensional wave operator $\partial_u \partial_v$. Secondly, the discrete variable can be made to correspond to the simple roots (Dynkin diagram) of a finite-dimensional or affine Lie algebra. The equation now looks like

$$\partial_u \partial_v \varphi_j = \exp - \sum_k C_{jk} \varphi_k, \quad (2)$$

where C_{jk} is the Cartan matrix of the Lie algebra [2–5]. In the original (infinite) Toda chain, the Lie algebra is the $N \rightarrow \infty$ limit of $A_{N-1} = su(N)$.

The subject of this note is the question of whether there exists an integrable version of (2) in which the variables u and v are discrete, i.e. an integrable partial difference equation which reduces to (2) in a continuum limit. The general topic of discrete integrable versions of partial differential equations has long been of interest. There have been several different approaches, for example:

- discrete version of the AKNS linear system [6];
- Hirota’s method [7–12];
- Zakharov-Shabat dressing method [13];
- linear integral equations [14–17].

In connection with discrete versions of the Lax system for the Toda chain (1), see also refs 18–21. Suris [18–20] exhibited discrete integrable versions of the Toda chain corresponding to all the non-exceptional affine Lie algebras. The question here is whether there are analogous integrable lattice versions of the Toda field equations (2). By “integrable” we mean that the equation can be expressed as the consistency condition of a linear system of the type described below. Two particular examples that have been known for some time are:

- a discrete sine-Gordon equation [9], corresponding to the algebra $A_1^{(1)}$;
- a discrete A_∞ Toda field equation [13, 12, 22].

The latter is often known as the Hirota or Hirota-Miwa equation. It is fairly straightforward to impose boundary conditions (in the index j) which reduce it to a discrete A_n or $A_n^{(1)}$ Toda field equation; but the resulting finite-rank systems do not seem to have appeared explicitly in the literature before. Let us therefore begin by presenting these, in a form similar to that of ref 18.

Let u and v be coordinates on the two-dimensional lattice $\mathbf{Z}^2$, with lattice spacing h . So each of u and v takes values which are integer multiples of h . A subscript denotes shift in the forward direction: in other words, if $f(u, v)$ is a function on the lattice, then $f_u(u, v) = f(u + h, v)$ and similarly for f_v . Consider a linear system involving $N \times N$ matrices U and V (with $N \geq 2$), and a column N -vector ψ , of the form

$$\psi_u = U\psi, \quad \psi_v = V\psi, \quad (3)$$

where

$$\begin{aligned} U &= \lambda^{-1}I + h \exp(F_u)X_- \exp(-F), \\ V &= \lambda \exp(F_v - F) - hX_+. \end{aligned} \quad (4)$$

The various symbols appearing in (4) are defined as follows:

- λ is an arbitrary scalar parameter;
- I is the identity matrix;
- $F = \text{diag}(f_1, f_2, \dots, f_N)$ is a diagonal matrix;
- X_+ is the matrix with ones immediately above the main diagonal, and zeros elsewhere, i.e. $(X_+)_{ab} = \delta_{a-b+1}$;
- X_- similarly has ones just below the main diagonal.

Note that X_+ is the sum of elements of A_{N-1} corresponding to its simple roots, in the usual representation. We are thinking in terms of a specific representation of the Lie algebra, because the discrete Toda equations are not purely Lie-algebraic (unlike in the continuum case).

The consistency condition for (3) is $U_v V = V_u U$. The λ^{-1} and λ terms of this equation are automatically satisfied by (4), and the remaining (λ -independent) term is equivalent to

$$\begin{aligned} \exp(\Delta^2 f_1) &= [1 + h^2 \exp(f_{2u} - f_{1v})], \\ \exp(\Delta^2 f_k) &= [1 + h^2 \exp(f_{(k+1)u} - f_{kv})] / [1 + h^2 \exp(f_{ku} - f_{(k-1)v})] \\ &\quad \text{for } 2 \leq k \leq N-1, \end{aligned} \quad (5)$$

$$\exp(\Delta^2 f_N) = 1 / [1 + h^2 \exp(f_{Nu} - f_{(N-1)v})],$$

where

$$\Delta^2 f = f_{uv} - f_u - f_v + f. \quad (6)$$

The equations (5) are the discrete A_{N-1} Toda field equations. Strictly speaking, we should impose the constraint

$$\sum_{k=1}^N f_k = 0, \quad (7)$$

so that there are only $N - 1$ independent fields. This is consistent with the field equations, because one of the equations (5) is in effect $\det(U_v V) = \det(V_u U)$, and this is automatically satisfied if (7) holds. So one should think of (5) as consisting of $N - 1$ second-order difference equations for $N - 1$ functions.

Let us examine two limiting cases of these field equations. First, if the f_k depend only on the single discrete variable $t = u + v$, then (5) reduces to the discrete A_{N-1} chain in the form described by Suris [18,19]. He does not impose the constraint (7), but remarks that the quantity $\Delta_t \sum f_k$ is a constant of motion. Secondly, in the continuum limit $h \rightarrow 0$ the equations (5) do indeed reduce to (2), where C_{jk} is the Cartan matrix of A_{N-1} and $\varphi_k = \sum_{j=1}^k f_j$ for $1 \leq k \leq N - 1$.

The equations corresponding to the affine algebra $A_{N-1}^{(1)}$ are very similar. The linear system is again (4), but now the constant matrix X_+ has an additional unit entry, in its lower left-hand corner: $(X_+)_{ab} = \delta_{a-b+1} + \delta_{a-N}\delta_{b-1}$. Similarly, X_- has a one in its upper right-hand corner. The resulting field equations are

$$\exp(\Delta^2 f_k) = [1 + h^2 \exp(f_{(k+1)u} - f_{kv})] / [1 + h^2 \exp(f_{ku} - f_{(k-1)v})] \quad (8)$$

for $1 \leq k \leq N$, where $f_0 = f_N$ and $f_{N+1} = f_1$ (in other words, f_k is periodic in the index k , with period N).

The two special cases described above for A_{N-1} work in exactly the same way for these $A_{N-1}^{(1)}$ equations. Another special case worth pointing out is that of $A_1^{(1)}$: taking $f_1 = -f_2 = i\varphi$, equation (8) with $N = 2$ reduces to the discrete sine-Gordon equation for φ [9].

By making different choices of the matrices F and $X_{\pm}$, Suris [18] obtained discrete Toda chains corresponding to other affine Lie algebras. Another way of deriving the same results is to use a symmetry reduction (cf. ref 5 for the continuum case). For the one-dimensional chain, where $f_k = f_k(t) = f_k(u + v)$, the equations (5) and (8) have the $\mathbf{Z}_2$

symmetry

$$f_N \mapsto -f_1, \quad f_{N-1} \mapsto -f_2, \quad \dots, \quad f_1 \mapsto -f_N. \quad (9)$$

So we can reduce by putting $f_N = -f_1$ etc. The A_{N-1} chain then reduces to the chain corresponding to $C_{N/2}$ (for N even) or $B_{(N-1)/2}$ (for N odd); and $A_{N-1}^{(1)}$ reduces to the chain $C_{N/2}^{(1)}$ (for N even) or $A_{(N-1)}^{(2)}$ (for N odd). Finally, the chain $D_{N/2}^{(2)}$ is obtained from $A_{N-1}^{(1)}$ with N even, via an analogous but slightly different reduction. Chains corresponding to other Lie algebras such as D_N involve a Lax representation of a more general type, and do not fit into the scheme presented here. An examination of some particular examples illustrates that these equations, despite the notation, are not strictly Lie-algebraic. For example, the discrete Toda chains labelled by A_1 and B_1 are different, even though these Lie algebras are isomorphic.

For the (two-dimensional) Toda field equations (5) and (8), however, things are not quite so simple. These equations do not admit the symmetry (9). But they do admit such a symmetry if we also interchange the independent variables u and v . In other words, we can reduce (5) and (8) by imposing

$$f_N(u, v) = -f_1(v, u), \quad f_{N-1}(u, v) = -f_2(v, u), \dots \quad (10)$$

Because of the nonlocal nature of this constraint, one then gets nonlocal field equations. But one can restore locality by restricting to symmetric functions of u and v . In effect, this yields a partial difference equation on the half-plane, rather than on the full plane $\mathbf{Z}^2$. Let us examine a particular example, namely the systems corresponding to B_n .

So let $N = 2n + 1$ be an odd integer, with $n \geq 1$. Impose (10), and also the condition that f_{kv} be symmetric in (u, v) , ie.

$$f_k(u, v + h) = f_k(v, u + h) \quad \text{for } k = 1, 2, \dots, n. \quad (11)$$

Finally, impose $f_{n+1} = 0$, which is consistent with (5, 10, 11). The effect of all this is that the field equations (5) reduce to

$$\begin{aligned} \exp(\Delta^2 f_1) &= [1 + h^2 \exp(f_{2u} - f_{1v})], \\ \exp(\Delta^2 f_k) &= [1 + h^2 \exp(f_{(k+1)u} - f_{kv})] / [1 + h^2 \exp(f_{ku} - f_{(k-1)v})] \\ &\quad \text{for } 2 \leq k \leq n-1, \\ \exp(\Delta^2 f_n) &= [1 + h^2 \exp(-f_{nv})] / [1 + h^2 \exp(f_{nu} - f_{(n-1)v})], \end{aligned} \quad (12)$$

together with the symmetry constraint (11). So we can think of (12) as being defined on the half-plane $v \leq u + h$. In terms of the laboratory coordinates $t = \frac{1}{2}(u + v - h)$ and $x = \frac{1}{2}(u - v + h)$, this half-plane consists of those lattice sites which lie in the region $x \geq 0$. The field equation is (12) at lattice sites for which $v < u + h$, and there is a boundary condition if $v = u + h$. The latter is obtained from (12) by using the symmetry about $x = 0$: for each k , f_{kv} is replaced by f_{ku} .

The continuum limit $h \rightarrow \infty$ of this gives the B_n Toda field equation in the half-plane $x \geq 0$, with the boundary condition $\partial_x \varphi_k = 0$ on $x = 0$. Notice also that the discrete B_n Toda chain is a special case, since the f_k are then independent of x .

In conclusion, it has been shown that there exist integrable discrete versions of the two-dimensional Toda field equations, corresponding to at least some of the simple finite-dimensional or affine Lie algebras. The basic cases are A_n and $A_n^{(1)}$. Others such as B_n and C_n can be obtained from these by reduction, but it appears that these reduced systems are defined on the half-lattice rather than the full two-dimensional lattice. Not all the reductions have as yet been analysed. And for the algebras D_n , $B_n^{(1)}$, $D_n^{(1)}$ and $A_{2n-1}^{(2)}$ the picture is unclear: the corresponding discrete Toda chains involve Lax representations of a more general type [18,19], and it is an open question whether discrete two-dimensional versions exist. Similarly, the situation with regard to the exceptional Lie algebras is unknown. It would be preferable to have a “group” description which did not involve a particular representation, but it is not clear what sort of algebraic structure is required for this.

References.

- [1] M. Toda, Theory of nonlinear lattices (Springer, 1988).
- [2] A. V. Mikhailov, M. A. Olshanetsky and A. M. Perelomov, Comm. Math. Phys. 79 (1981) 473.
- [3] G. Wilson, Ergod. Th. and Dynam. Syst. 1 (1981) 361.
- [4] A. P. Fordy and J. Gibbons, Proc. R. Ir. Acad. A 83 (1983) 33.
- [5] D. Olive and N. Turok, Nucl. Phys B 215 (1983) 470.
- [6] M. J. Ablowitz and F. J. Ladik, Stud. Appl. Math. 55 (1976) 213; 57 (1977) 1.
- [7] R. Hirota, J. Phys. Soc. Jpn. 43 (1977) 1424.
- [8] R. Hirota, J. Phys. Soc. Jpn. 43 (1977) 2074.
- [9] R. Hirota, J. Phys. Soc. Jpn. 43 (1977) 2079.
- [10] R. Hirota, J. Phys. Soc. Jpn. 45 (1978) 321.
- [11] R. Hirota, J. Phys. Soc. Jpn. 46 (1979) 312.
- [12] R. Hirota, J. Phys. Soc. Jpn. 50 (1981) 3785.
- [13] D. Levi, L. Pilloni and P. M. Santini, J. Phys. A 14 (1981) 1567.
- [14] F. W. Nijhoff, G. R. W. Quispel and H. W. Capel, Phys. Lett. A 97 (1983) 125.
- [15] F. W. Nijhoff, H. W. Capel, G. L. Wiersma and G. R. W. Quispel, Phys. Lett. A 103 (1984) 293; 105 (1984) 267.
- [16] G. R. W. Quispel, F. W. Nijhoff, H. W. Capel and J. van der Linden, Physica A 125 (1984) 344.
- [17] F. W. Nijhoff, V. G. Papageorgiou and H. W. Capel, Springer Lecture Notes in Mathematics 1510 (1992) 312.
- [18] Yu. B. Suris, Leningrad Math. J. 2 (1990) 339.
- [19] Yu. B. Suris, Phys. Lett. A 145 (1990) 113.
- [20] Yu. B. Suris, Phys. Lett. A 156 (1991) 467.
- [21] J. Gibbons and B. A. Kupershmidt, Phys. Lett. A 165 (1992) 105.
- [22] S. Saito and N. Saitoh, J. Math. Phys. 28 (1987) 1052.